## Abstract Submitted to the International Conference on Strongly Correlated Electron Systems University of Michigan, Ann Arbor August 6-10, 2001

## $\mu$ SR studies of unconventional superconductivity in Ce TIn<sub>5</sub> (T=Ir, Co)

Wataru Higemoto<sup>1</sup>, Akihiro Koda<sup>1</sup>, Ryosuke Kadono<sup>1</sup>, Yoshinori Haga<sup>2</sup>, Dai Aoki<sup>3</sup>, Hiroaki Shishido<sup>3</sup>, Rikio Settai<sup>3</sup>, Yoshichika Ōnuki<sup>2,3</sup>

- Meson Science Laboratory, Institute of Materials Structure Science, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan
- <sup>2</sup> Advanced Science Research Center, Japan Atomic Energy Research Institute, Tokai, Ibaraki 319-11, JAPAN
- <sup>3</sup> Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043

We report on muon spin rotation/relaxation ( $\mu$ SR) measurements in heavy fermion superconductors CeIrIn $_5$  and CeCoIn $_5$  to elucidate the origin of unconventional superconductivity through their magnetic properties.

In CeIrIn<sub>5</sub>, we demonstrate the absence of static long range magnetic ordering down to 20mK under zero magnetic field. In a magnetic field of 400G ( $>H_{c1}$ ), the muon spin relaxation due to the inhomogeneous magnetic induction of flux line lattice sets in below 0.4K, showing that bulk superconductivity occurs around 0.4K while the zero resistance is observed above 1 K. The spin relaxation rate shows gradual increase with decreasing temperature down to 20mK, indicating that the quasiparticle excitation does not fall exponentially with temperature. This result strongly suggests the superconductivity with non-s wave pairing in this compound. The result of  $\mu$ SR study on CeCoIn<sub>5</sub> will be also reported.